

**PORCUPINE CARIBOU
INTERIM ANNUAL SUMMARY REPORT
2020 – 2021**



Barn Mountains, May 2020

Submitted to: Porcupine Caribou Management Board

Submitted by: Porcupine Caribou Technical Committee

December 9, 2021

Indicator Table

Annual Summary Report 2020 – 2021

Indicator	Value	Average	Notes
Population size and trend			
Population size	2021 = photocensus not attempted	--	Herd did not aggregate sufficiently for photocensus. Successful photocensus in 2017 found 218,457 (95% CI = 202,106 - 234,808) caribou.
Population trend	2021 = no data	$\Lambda = 1.037$ (2010-2017)	Declined by 55,000 caribou between 1989 and 2001. Recovered to 169,000 by 2010 and continued to increase to 218,457 in 2017. Annual growth rate almost identical to growth phase from 1972-1989.
Adult cow survival	2019 = 90.5 % (95% CI = 80.1 – 95.6%)	88.2 % (2012-2019)	Sampling interval is June 1 2020 – July 30 2021. Data for this past year will be provided before the AHM.
Adult bull survival	2019 = 73.4 % (95% CI = 44.1 – 89.1%)	70.1 % (2015 – 2019)	Sampling interval is June 1 2020 – July 30 2021. Data for this past year will be provided before the AHM.
Yearling cow survival	2019 = 86.3 % (95% CI = 55.8 – 96.4%)	86.7 % (2017 – 2019)	Sampling interval is June 1 2020 – July 30 2021. Data for this past year will be provided before the AHM.
Calf birth rate (Parturition rate)	≥ 4 yr olds = 0.85	0.85 (5-yr average)	35-year average = 0.82
	≥ 3 yr olds = 0.83	0.82 (5-yr average)	15-year average = 0.79
	3 yr olds = 0.71	--	Small sample sizes for each year and limited consecutive years limit the ability to calculate meaningful averages or adequate time-series weighted averages.
Post-calving survival	2021 = 0.90	0.87 (long- term average)	Includes ≥ 3 year old adult cows
Late June calf:cow ratio	2021 = 0.63	0.57 (long- term average)	Includes ≥ 3 year old adult cows
March calf:cow ratio	2021 = no data	--	17-year average = 0.35. Survey in March 2017 = 0.35
Bull Ratio	2021 = no data	--	
Peak of calving	2021 = May 30	31 May	Peak of calving in 2021 was May 30, slightly earlier than average

Indicator Table

Annual Summary Report 2020 – 2021

Body condition			
Average backfat	F: 1.3 cm; M: 2.6 cm	F: 2.6 cm; M: 2.0 cm	Caribou generally not available for fall 2020 harvest, majority of samples collected November 2020 thru March 2021
Hunter assessment	F: 2.8 M: 3.0	F: 2.2 M: 3.0	Caribou generally not available for fall 2020 harvest, majority of samples collected November 2020 thru March 2021
Overall condition of caribou	2020-21 = generally good	Good	Caribou Body Condition generally remained good
Habitat and other considerations			
Snow conditions winter (2020-21)	<u>Snow Depth</u> Eagle = 72.7 cm Ogilvie = 71.3 cm Old Crow = 63.0 cm North Slope = no data Richardson = no data <u>Snow Density</u> Eagle = 0.18 g/cm ³ Ogilvie = 0.20 g/cm ³ Old Crow = 0.17 g/cm ³ Richardson = no data North Slope = no data	<u>5 yr Av. Depth</u> 76.9 cm 65.8 cm 70.0 cm no data no data <u>Density</u> 0.19 g/cm ³ 0.18 g/cm ³ 0.18 g/cm ³ no data no data	Eagle Region was above long term average, but below the 5 year average; Ogilvie Region was above both the long and 5 year average; Old Crow Region was slightly above the long term average, but below the 5 year average. Snow density in all areas were slightly below the 5 year average, and below the long term average in all areas but Ogilvie. Eagle and Old Crow Regions appear to have an increasing trend in snowpack.
Wildland fires	2021 = data not yet available 2020 = 211.5 km ²	5 year average 1551.9 km ²	No significant burns in 2020. 1 large fire occurred south of the Porcupine River and several on the periphery of the herds range in the Peel drainage during 2021.
Linear disturbance and human development	2020-21 = No major increases	N/A	Oil and gas lease sales proposed for 2021 for the 1002 lands in Alaska. Seismic work proposed for a portion of 1002 east of Marsh Creek from January-September 2021. Some oil and gas drilling may be proposed for the Eagle Plains area in 2021.

Contents

Section	page
Indicator Table.....	2
Indicator Table.....	3
INTRODUCTION.....	7
Porcupine Caribou Annual Summary Report.....	7
Herd background.....	7
Management direction and goals.....	7
POPULATION.....	8
Population size – photocensus.....	8
Population size – computer modeling.....	10
Survival Estimates.....	12
Calf birth rate and calf survival.....	14
Peak and Extent of calving grounds.....	18
Table 2. Peak dates of calving for the Porcupine Caribou herd.....	19
Bull Ratio.....	19
CARIBOU BODY CONDITION.....	20
Hunter assessments and condition indicators.....	20
HABITAT.....	23
Wildland fires.....	23
Linear disturbance and human development footprint.....	26
Snow condition.....	28
LITERATURE CITED.....	31
Appendix A. Summary of biological parameters.....	33
Appendix B. Previous research findings.....	35
Adult female survival.....	35
Short yearling survival to 3 years of age.....	36
Adult bull survival.....	37

	page
Figure 1. Map of the Porcupine caribou herd from June 28 to July 9 th , 2021 showing large easterly movements that did not provided ideal aggregations to conduct a photocensus.	9
Figure 2. Population size of the PCH from 1972 to 2017. Data from 1972 to 2001 are photocensus minimum counts. Estimated abundance and associated 95% confidence intervals in 2010, 2013 and 2017 were derived from photocensus minimum counts and modeling (Rivest et al. 1998).	10
Figure 3. Projected population size of the Porcupine caribou herd under five different calf mortality scenarios (e.g., 0.4 represents 40% of calves dying or 60% surviving from birth to 1 year of age; note this is the highest increase of the five scenarios presented while 0.58 or 58% mortality would equate to the lowest). The grey represents the uncertainty around the five different scenarios with darker shades of grey representing greater overlap and a more likely outcome.	11
Figure 4. Year specific survival estimates for GPS collared adult female Porcupine caribou (black circles) compared to the grand mean (solid horizontal bar), 2012–2019 (Caikoski 2020b). All confidence intervals are 95%. A year is defined as the period from caribou birth (June 1) to one year later (May 31). For example, year 2012 occurs from June 1, 2012–May 31, 2013.	13
Figure 5. Locations of collared Porcupine caribou cows observed with a newborn calf (green circles), those judged to be parturient but had not yet given birth (blue squares), and those judged to be barren (black triangles), 29 May–10 June 2021. Extent of the calving grounds was estimated by the isopleth encompassing 99% of the fixed kernel utilization distribution of locations of cows observed with a calf (purple polygon). The concentrated calving area is the area with greater than average density of caribou cows with calves (red polygon). (Caikoski 2021)	16
Figure 6. Estimated birth rate, calf survival indices and March composition count for the Porcupine Caribou herd from 1985-2021.	17
Figure 7. Locations of GPS collared Porcupine caribou cows observed with a calf (green circles) and without a calf (red circles) during 19–20 June 2021. (Caikoski 2021). ..	18
Figure 8. Average condition of harvested Porcupine caribou recorded by hunters. 1=poor 2=fair 3=good 4=very good. Error bars are standard errors. Labels indicate # of caribou sampled.	22
Figure 9. Average depth of backfat (cm) recorded in Body Condition Monitoring. Error bars are standard errors. Labels indicate # of caribou sampled.	22
Figure 10. Areas burned within range of the Porcupine Caribou Herd in Yukon, NWT and Alaska from 1960 to 2020.	24
Figure 11. Total number of fires and number of large fires to 2020 within the range of the Porcupine Caribou Herd in Yukon, NWT and Alaska.	25
Figure 12. Total area burned by fire, by year to 2020 within the range of the Porcupine Caribou Herd in Yukon, NWT and Alaska.	25
Figure 13. Human disturbance within the range of the Porcupine Caribou herd (updated Nov. 14, 2014 – November 23, 2017) in Alaska, Northwest Territories, and Yukon.	27
Figure 14. The extent of 3D seismic lines, trails and roads cut or brushed out in 2013-14 by Northern Cross within the range of the Porcupine Caribou Herd in Yukon	27
Figure 15. Winter distribution of Porcupine Caribou from Dec. 1, 20120 to March 31, 2021.	28

Figure 16. Summary of snow depth and density by snow region from permanent stations (indicated by green stars) for the Yukon portion of the Porcupine Caribou Herd range. Red lines on the map delineate snow regions relevant to caribou (Russell et al 1993). 30

Figure 17. Annual survival estimates for adult female Porcupine Caribou, May 2003 – June 2012. Source: USFWS unpublished data. 36

Figure 18. Survival of Porcupine Caribou females from 9 months to 3 years of age from 2003-2010. 37

Figure 19. Survival of male Porcupine Caribou from 2003 to 2010. 38

List of Tables

	page
Table 1. Estimates of annual survival for Porcupine Caribou Herd adult females (2012–2019), adult males (2015–2019), and yearling females (2017–2019; Caikoski 2020b)..	14
Table 2. Peak dates of calving for the Porcupine Caribou herd.....	19

INTRODUCTION

Porcupine Caribou Annual Summary Report

This report was prepared for the Porcupine Caribou Management Board (PCMB) to provide information to make an assessment on the status of the herd as part of the *Harvest Management Plan for the Porcupine Caribou Herd in Canada* (HMP). Information within this report was guided by the topics listed in the HMP. As noted in relevant sections, some information is not available or analyzed. Under the HMP, Parties are requested to comment on this report and provide additional information to the PCMB at the Annual Harvest Meeting. Information for this summary report was provided by members of the Porcupine Caribou Technical Committee (PCTC) to inform that process.

Herd background

The Porcupine Caribou Herd's (PCH) core home range is approximately 201,190 km² and extends into Alaska, Yukon, and the Northwest Territories. Within this range there are currently 12 different areas where different agencies have jurisdiction over land and/or wildlife management. Management of the herd must take into consideration:

- 2 federal governments
- 3 state or territorial governments
- 8 Indigenous land claim agreements
- 5 national parks or preserves
- 1 territorial park
- 2 special management areas
- 2 specific ordinances
 - Dempster Highway Area Development Ordinance, and
 - a federal Order-in-Council Withdrawal (Yukon North Slope)

The PCH was the first international caribou herd with its own formal co-management agreements and boards. There are five main management agencies which work on the herd: Canadian Wildlife Service, U.S. Fish and Wildlife Service, Government of Yukon, Government of the Northwest Territories, and the Alaska Department of Fish and Game. Management and research is coordinated by the PCTC which consists of biologists from numerous agencies, co-management boards as well as occasional faculty members or students from various universities.

All Indigenous organizations within the Canadian range of the herd have land claim agreements. These agreements solidify the Indigenous right to hunt for subsistence and ensure local participation in wildlife management through co-management boards. The agreements also created lands that are privately owned and managed by the First Nations or Inuvialuit. Self-governing agreements in Yukon also give the First Nation governments the ability to regulate their citizens and their land.

Management direction and goals

To help coordinate management, two Porcupine Caribou agreements were set up, each creating a co-management board. In 1985, three governments and three Indigenous organizations signed the *Porcupine Caribou Management Agreement* (PCMA), creating the within-Canada Porcupine Caribou Management Board. In 1987, an *Agreement Between the Government of Canada and the Government of United States on the Conservation of the Porcupine Caribou Herd* was signed, creating the International Porcupine Caribou Board (IPCB).

The PCTC drafts work plans to coordinate research and monitoring activities, optimize funds and staff time, and provide technical information to co-management boards and agencies. Harvest

management is co-operative among the Parties to the PCMA and is guided by the HMP and the accompanying Implementation Plan.

Goals that pertain to the PCTC taken from the *Porcupine Caribou Management Agreement* (1985) and reflected in subsequent documents prepared by the PCMB include:

B. The Board shall review relevant scientific information [and traditional knowledge] on the conservation management of the herd and its habitat, and make recommendations to the Minister on policy, legislation and regulations regarding:

- Management strategies
- Further research where there appears to be a need, including recommendations on methods of data collection and presentation;
- A herd management plan; and
- A predator management plan.

D. The Board may identify sensitive [caribou] habitat areas requiring special protection and recommend measures to protect such areas.

The *Plan for the International Conservation of the Porcupine Caribou Herd* outline a number of objectives pertinent to the PCTC.

- To conserve the Porcupine caribou herd and its habitat through international cooperation and coordination so that the risk of irreversible damage or long-term adverse effects as a result of use of caribou or their habitat is minimized.
- To ensure opportunities for customary and traditional uses of the Porcupine caribou herd.
- To enable users of Porcupine caribou to participate in the international coordination of the conservation of the Porcupine caribou herd and its habitat.
- To encourage cooperation and communication among governments, users of Porcupine caribou, and others to achieve the objectives of the Agreement.

POPULATION

Population size – photocensus

Objective

To estimate the size of the herd every 2 to 3 years.

Methods

A technique called an Aerial Photo Direct Count Extrapolation has been used to estimate the herd size since 1972 (Davis 1979, Valkenburg et al. 1985, Rivest et al. 1998). Once the insects come out during the warm weather in late June or early July, the caribou gather into very large, tight groups sometimes consisting of tens of thousands of caribou. These large groups are photographed and caribou in the photos are counted. Any caribou that are found outside of the large groups are added and the estimate is rounded to the nearest thousand caribou. Radio-collared caribou are used to help locate the caribou aggregations and correct the estimate for any missing caribou. This technique is considered an accurate and reliable method to count large barren-ground caribou herds and can also provide a measure of uncertainty (confidence interval) around the population estimate. A confidence interval is a range of values that describes the uncertainty surrounding the population estimate. For example, the photocensus in 2013 found that the population estimate of the PCH was 197,228 (95% CI = 168,667 – 225,789). That means that we are 95% confident that the true population estimate is within the upper (225,789) and lower number (168,667).

Results

The herd was monitored using GPS collar locations throughout June and July 2020 to see if the herd would form large aggregations to allow for a photocensus. In late June the herd initiated an eastern movement that caused caribou to be more dispersed than what is required for a photocensus attempt (Figure 1. Map of the Porcupine caribou herd from June 28 to July 9th, 2021 showing large easterly movements that did not provided ideal aggregations to conduct a photocensus.

The last successful photocensus was completed in 2017 and resulted in a minimum count of 198,104 caribou and a population abundance estimate of 218,457 (95% CI = 202,106 – 234,808) caribou (Caikoski 2017). The annual growth rate from 2010 to 2017 was estimated at $\lambda = 1.037$ (SE = 0.0082; Caikoski 2017).




Figure 1. Map of the Porcupine caribou herd from June 28 to July 9th, 2021 showing large easterly movements that did not provided ideal aggregations to conduct a photocensus. Note, some collared caribou used to produce this map may be considered as part of the Central Arctic Herd (specifically the northwest North Slope polygon data) and require further review. Data will be reassessed at a later time and this map may need to be revised to most accurately reflect PCH distribution over this period.

Discussion

The most recent photocensus in 2017 indicated that the Porcupine Caribou herd was at its highest number since surveys began in the 1970's (Figure 2). Analysis of how fast the herd is

growing shows that the current growth rate ($\lambda = 1.037$: Caikoski 2017) is almost identical to the growth rate during the last growth phase of the herd from 1972 to 1989. If this number ($\lambda = \text{lambda}$) is larger than 1 then population is increasing. A population with a $\lambda < 1$ may indicate a decreasing population. A photocensus effort is planned for summer 2021.




Figure 2. Population size of the PCH from 1972 to 2017. Data from 1972 to 2001 are photocensus minimum counts. Estimated abundance and associated 95% confidence intervals in 2010, 2013 and 2017 were derived from photocensus minimum counts and modeling (Rivest et al. 1998).

Population size – computer modeling

Objective

To estimate the total herd size and the measure of uncertainty surrounding that estimate using a computer population model. The model is also useful for examining the influence of some of the different herd indicators (e.g., adult female survival or the number of calves) and the effect when you combine each of these within a year. Uncertainty surrounding the actual size of the population increases as the model projects further into the future from the last known photocensus (e.g., 2017).

Methods

Variables in the model include: previous photocensus estimates, and annual estimates of adult sex ratio, harvest numbers, cow survival, and calf recruitment. We also included a measure of how certain we are about these values into the model. This allows us to have a measure of uncertainty in our final herd size estimate(s) (i.e., confidence interval). We calculated the uncertainty in the final model projections by running the model many times, each time using a different combination of these variable estimates. We do this because we never know the precise

number of calves or exactly how many cows die in a year, but we know approximately what these numbers are from our monitoring. The uncertainty provided around projections is useful, as it tells us how confident we are in the size and trend of the population estimate and the probability that we are in a specific management zone identified by the HMP.

To develop the model in 2020, we used information from the indicators collected (see Annual Summary Reports) since the last successful photocensus (i.e., 2017). We modelled survival of different age and sex classes of the herd using available information where possible (e.g., yearling cows, adult males and females). When information was not available (yearling, 2-3 year old bulls), we estimated these rates based on the most similar age and sex groupings that had data and adjusted according to available information (e.g., yearling bull survival is lower than yearling cows but higher than adult bulls). Where information was only partially available (e.g., total harvest each year), we used the best available information and estimated the actual harvest for the year to ensure each Party had at least some harvest accounted for. For calf recruitment, we did not have any data available other than parturition rates and late June survival, however field staff do keep qualitative track of the number of calves throughout the range during capture events. We used this information together with historic recruitment data and ran five different recruitment scenarios in our population projections to get a general sense of the population and trend. We chose to use a range of values (0.4 – 0.58) to represent calf mortality rate (i.e., recruitment) as it was the variable that we lack sufficient data for.

Results

We produced five different trend lines with confidence intervals surrounding each line for 2017-2020 (Figure 3). All scenarios and their associated confidence intervals fell within the threshold for the Green Zone as identified by the HMP (i.e., >115,000 caribou). Most scenarios indicate that the herd has remained either stable or possibly increased, although a proportion of runs where calf survival was extremely poor shows the herd may have declined slightly.




Figure 3. Projected population size of the Porcupine caribou herd under five different calf mortality scenarios (e.g., 0.4 represents 40% of calves dying or 60% surviving from birth to 1 year of age; note this is the highest increase of the five scenarios presented while 0.58 or 58% mortality would

equate to the lowest). The grey represents the uncertainty around the five different scenarios with darker shades of grey representing greater overlap and a more likely outcome.

Discussion

Given the difficulty with obtaining regular photocensus estimates of the herd's size, primarily due to poor weather or a lack of animal aggregation, this tool was designed to provide the Board and Parties with information on the herd's population size and trend. The population model incorporates uncertainty, which is critical information when managers are developing management actions.

Working from the 2017 population estimate of 218,457 (95% CI = 202,106 – 234,808) the models project that the herd is likely stable or possibly increasing, with a smaller chance of a decline. Based on these results it is extremely likely that the herd remains in the Green Zone.

To generate the models several assumptions are required. Harvest data is available in part each year, however, precise estimates of total harvest and the sex ratio of harvest remain coarse estimates. Having more precise harvest estimates wouldn't likely change the population estimate, but improved harvest data would result in a better estimate with less uncertainty. The addition of annual survival estimates for adult cows and bulls, and for yearling cows, completed by the PCTC, is a significant improvement on historic survival estimates that were limited based on sample size, duration of each study, and the required analytical framework used. The largest gaps in data at this time appear to be calf recruitment, the sex ratio of the herd, and survival of younger bull classes.

While modelling all of the information we collect from monitoring can be useful in understanding the trajectory and the amount of growth or decline that may be occurring, it should be noted that caution is needed, particularly as we move further from the 2017 population estimate. Even small amounts of error in some sources of data, though addressed in our modelling approach, can still lead to reduced accuracy of models. Compounding that error year over year means our confidence becomes reduced each year, as demonstrated by the growing area in grey found in Figure 3. Using all sources of knowledge together will yield the best understanding of the status of the herd.

Survival Estimates

Objective

To obtain an annual estimate of survival for adult female, adult male, and yearling female Porcupine caribou.

Methods

Annual survival was estimated from GPS collared caribou using known-fate models (logistic regression). Annual survival for adult females (years 2012–2019), adult males (years 2015–2019), and yearling females (years 2017–2019) were conducted separately and were reported with 95% confidence intervals. A year was defined as June 1 through May 31, which represents the time period from birth to consecutive birth dates.

Results

The sampling period was from June 1, 2019 through May 31, 2020. Adult cow survival for this time period was estimated to be 90.5 % (95% CI = 80.1 – 95.6%), slightly above the average of 88.2 % (2012-2019). Adult bull survival was 73.4 % (95% CI = 44.1 – 89.1%), also slightly above the average of 70.1 % (2015 – 2019). Yearling cow survival was 86.3 % (95% CI = 55.8 – 96.4%), on par with the average of 86.7 % (2017 – 2019).

From 2012–2019, annual survival of adult females ranged from 78.5 % (95% CI = 63.9 – 87.8%) to 94.5% (95% CI = 83.9 – 98.2%) and averaged 88.2% (95% CI = 84.3 – 91.1%) across all years (Caikoski 2020b; Table 1. Estimates of annual survival for Porcupine Caribou Herd adult females (2012–2019), adult males (2015–2019), and yearling females (2017–2019; Caikoski 2020b). A year is defined as June 1-May 31 (e.g. year 2012 = June 1 2012–May 31 2013). Figure 4).

During 2015–2019, annual survival of adult males ranged from 58.7% (95% CI = 19.8 –84.3%) to 73.4% (95% CI = 44.1 – 89.1%) and averaged 70.1% (95% CI = 58.3 –79.1%; Caikoski 2020b; Table 1).

During 2017–2019, annual survival of yearling females ranged from 80.8% (95% CI = 51.9% – 93.4%) to 93.2% (95% CI = 61.2% – 99.0%).




Figure 4. Year specific survival estimates for GPS collared adult female Porcupine caribou (black circles) compared to the grand mean (solid horizontal bar), 2012–2019 (Caikoski 2020b). All confidence intervals are 95%. A year is defined as the period from caribou birth (June 1) to one year later (May 31). For example, year 2012 occurs from June 1, 2012–May 31, 2013.

Discussion

Prior to the deployment of significant numbers of GPS collars on the herd, previous studies estimated survival rates of adult females using periodic radiotracking flights of VHF collars throughout the year and staggered entry product-limit methods (Kaplan and Meier 1958, Pollock et al. 1989). Fancy et al. (1994) reported an average annual survival rate of 84% during 1982–1991, Arthur et al. (2003) reported an average annual survival rate of 81% during 1997–2001, and Wertz et al. (2007b) reported an average annual survival rate of 82% during 2003–2006. Average annual survival of 88% during 2012–2017, suggests that survival of adult females improved compared to previous studies and is consistent with population growth during the same time period.

Population models based on PCH demographics suggest that relatively small but persistent reductions in adult female survival would result in population decline (Walsh et al. 1995, Griffith et al. 2002, Arthur et al. 2003). However, a suite of demographic parameters likely confound the effect of adult survival on abundance and can either mask the effects of high adult survival or may mitigate against poor adult survival. Furthermore, precision associated with estimates of survival to date are insufficient to detect statistical differences when small changes in vital rates occur. However, empirical evidence from vital rates reported here and three other studies compared to population abundance over the same time periods suggests a minimum long term average of 84% in annual survival for adult females may be necessary to prevent population decline.

Future estimates of yearling female survival will likely improve our understanding of recruitment to the 2-year-old age class, particularly during stable, growing, or declining phases in herd abundance.

Table 1. Estimates of annual survival for Porcupine Caribou Herd adult females (2012–2019), adult males (2015–2019), and yearling females (2017–2019; Caikoski 2020b). A year is defined as June 1–May 31 (e.g. year 2012 = June 1 2012–May 31 2013).

Age/Sex	Year	Annual Survival	95% LCL	95% UCL
Adult Females	2012	0.887	0.621	0.971
	2013	0.871	0.653	0.957
	2014	0.886	0.724	0.956
	2015	0.785	0.639	0.878
	2016	0.891	0.758	0.953
	2017	0.945	0.839	0.982
	2018	0.869	0.764	0.956
	2019	0.905	0.801	0.956
	Grand Mean	0.882	0.843	0.911
Adult Males	2015	0.717	0.415	0.883
	2016	0.693	0.444	0.848
	2017	0.716	0.451	0.871
	2018	0.587	0.198	0.843
	2019	0.734	0.441	0.891
		Grand Mean	0.701	0.583
Yearling Females	2017	0.932	0.612	0.990
	2018	0.808	0.519	0.934
	2019	0.863	0.558	0.964
		Grand Mean	0.867	0.728

Calf birth rate and calf survival

Objective

To document the annual calf birth rate and survival rate.

Methods

Calving surveys are conducted each year to estimate the birth rate and early survival rate of calves. Radio collared females ≥ 3 -years old are located from a fixed-wing aircraft and are classified as barren, pregnant, or have given birth. Researchers can tell if female caribou are pregnant if cows are observed with calves, have hard antlers or distended udders (Whitten 1995). Starting in 2018 we will be provided parturition rates for both 3-year old and ≥ 4 year old cows. Researchers found that 3-year old parturition rates might be useful as an index of nutritional status on the calving and summer ranges reflecting herd recovery or potential overgrazing issues with populations increases (Boertje et al. 2012). When 3-year-old parturition rates fall below $<55\%$ (5-year moving weighted average) this signals low nutrition due to potential overgrazing or adverse weather conditions.

Female caribou are re-located after about one month to determine whether the calves have survived. Calving success is described as the percent of cows that had calves. The July calf ratio is based on the proportion of collared females still with calves in late June or early July.

During the March composition count we do not use the ratio of collared females in late winter because the majority of calves will have weaned by March, but instead estimate the number of calves for every 100 adult cows, called a calf:cow ratio.

Results

Parturition rate

Staff from ADF&G conducted a survey from 29 May through 10 June, 2021 to estimate how many females were pregnant (parturition rate) in the population, though radiotracking flights did not occur on 31 May– 1 June, 5 June, or 8–9 June due to fog extending to the ground and poor visibility (Figure 5; Caikoski 2021).

The parturition rate for adult cows ≥ 4 -years of age and for adult cows ≥ 3 -years of age was slightly above average (85%, $n = 39$; 83%, $n = 46$) Figure 6). Parturition rate for 3-year-old cows was average at 71% ($n = 7$). We do not have a long enough data in the time-series to calculate 5-year weighted moving average for 3-year-old parturition rate to determine if prolonged low nutritional status may signal a declining population.

Post-calving survey

ADF&G staff conducted a post calving survey on 24 – 25 June, 2020 to estimate cow:calf ratio and mid-summer calf survival 3 weeks after peak of calving (Figure 7; Caikoski 2019a). Post-calving survival, estimated from cows observed with calves in early June that were subsequently observed in late June (excludes mortality shortly before and after birth), was 94% for calves of adults cows ≥ 3 -years of age ($n = 29$; Figure 6). The late June calf:cow ratio was 56 calves per 100 cows ≥ 3 -years of age or older ($n = 57$; Figure 6). This rate was identical to the long-term mean. During the survey, most post-calving caribou were concentrated on the Coastal Plain between Katakaturuk and Okpilak Rivers in Alaska (Figure 7).

March composition count

No composition count was conducted in March 2021. The last composition count (2017) reported a calf:cow ratio close to the long term average for the herd (35.8 calves per 100 cows; Figure 6).

Discussion

Population dynamics are most affected by survival of adult females over the medium and long term but can withstand fairly large annual fluctuations in calf birth rate or calf survival over the

short term. Figure 6 shows large fluctuations in these rates, but if birth rates or calf survival rates are low for several years in a row, population growth is more vulnerable.

Since 2017, we have documented declines in birth rate for all age classes and similar declines in late June calf:cow ratio. Although these measures have declined, birth rates (with the exception of ≥ 3 -year olds) and late-June survival are similar to long-term averages.

Three-year-old parturition rate is used as a long-term index that may reflect the impact of weather or range conditions on nutritional status. Based on past herd case studies, Boertje et al. (2012) suggest that managers can use this index to predict impending declines in herd numbers. We will continue to collect this information and compare to the threshold suggested by Boertje et al. (2012).




Figure 5. Locations of collared Porcupine caribou cows observed with a newborn calf (green circles), those judged to be parturient but had not yet given birth (blue squares), and those judged to be barren (black triangles), 29 May–10 June 2021. Extent of the calving grounds was estimated by the isopleth encompassing 99% of the fixed kernel utilization distribution of locations of cows observed with a calf (purple polygon). The concentrated calving area is the area with greater than average density of caribou cows with calves (red polygon; Caikoski 2021).




Figure 6. Estimated birth rate, calf survival indices and March composition count for the Porcupine Caribou herd from 1985-2021.




Figure 7. Locations of GPS collared Porcupine caribou cows observed with a calf (green circles) and without a calf (red circles) during 19–20 June 2021 (Caikoski 2021).

Peak and Extent of calving grounds

Objective

To estimate the date when greater than half of the collared adult female caribou have given birth each spring and the extent of the calving grounds.

Methods

During the calving surveys to document the birth rate (see previous), researchers record the date of their flights and how many of the collared cows have given birth. Only adult female caribou aged 3 years or older are used for this indicator. In some cases, the birth date is estimated based on the estimated age of the calf. The researchers then estimate the date when half of the collared adult female caribou have given birth. Peak of calving was approximated as the date at which greater than half of parturient cows were observed with a calf at heel.

Results

Of the cows ≥ 3 -years of age that were judged to be parturient, 18 of 31 were observed with a calf at heel by 1 June, indicating peak of calving occurred on May 30 (Caikoski 2021). Calving mostly occurred in the foothills of the Brook's Range between the Aichilik River and Katakaturuk River in Alaska, however some calving occurred on the coastal plain (Figure 4). Concentrated calving occurred in foothills along the Okpilak River drainage, Alaska.

Discussion

Caribou typically give birth *en masse* with many of the cows giving birth within days of each other. This is thought to be a strategy to reduce the risk of predation on any individual calf. This means that most of the cows would have been bred within a very short time period therefore peak of calving can be used as an indicator of how the rut went the previous fall. If the calving period is extended, it might mean that the rut was disrupted and cows were bred in a second estrus. This shows up as calves being born over an extended period of time. This is important because calves born late in the season are probably more likely to die from predators and they also may be too small to make the migration south for winter, reducing calf survival.

Since 1999, the peak date of calving has varied by a few days each year unlike some herds in central Canada that have seen a progressively later peak of calving. In 2021, peak of calving was on or slightly before the average date (June 2nd). We will continue to monitor this indicator to see if this is an emerging trend.

Table 2. Peak dates of calving for the Porcupine Caribou herd.

Year	Peak of calving	Note
1999	3-Jun	1 to 5 June
2000	7-Jun	
2001	8-Jun	5 to 10 June
2002	5-Jun	
2003	1-Jun	
2004	3-Jun	3 or 4 June
2005	2-Jun	1 to 4 June
2006	2-Jun	
2007	30-May	
2008	30-May	29 or 30 May
2009	2-Jun	Before 2 Jun
2010	2-Jun	
2011	2-Jun	
2012	No data	
2013	4-Jun	3 rd or 4 th June
2014	No data	
2015	No data	
2016	1-Jun	
2017	3-Jun	
2018	6-Jun	Could be bias late due to poor weather early in surveys
2019	4-Jun	
2020	1-Jun	
2021	30-May	
Average	2-June	

Bull Ratio

Objective

To document the ratio of bulls to cows in the herd.

Methods

We fly by helicopter during mid-October and classify as many as 200 caribou around each radio-collared caribou (bulls, cows, short yearlings). Caribou are classified into cow, calf, or either small, medium, or large bull. Then the number of bulls relative to the number of cows is calculated by dividing the total number of bulls by the total number of cows.

Results

Surveys are planned to coincide with years where photocensus surveys are completed, therefore surveys were not completed for 2020. The last survey was attempted by ADF&G in October 2017 but was unsuccessful due to poor weather.

The ratio of bulls to cows was estimated first in 1980 (Porcupine Caribou Management Plan 1989). That study estimated that there were about 60 bulls for every 100 cows which indicated a healthy herd. Bull survival and the bull ratio were not regularly monitored in following years because as long as the pregnancy rate remained high, there was no reason to believe that there are too few bulls to breed the cows. Subsequent surveys occurred in 2009 and 2010. Results from the 2010 survey are the most reliable and indicated a ratio of 57:100. In 2013 due to the poor result achieved in 2012 and the successful completion of a photo count on the herd, a rut survey was planned. Unfortunately leading up to the survey a large proportion of the herd moved to the western edge of the herd's range, eventually mixing with members of the Central Arctic Herd (CAH). Monitoring during the rut showed most caribou remained mixed with the CAH which resulted in the cancellation of the rut count.

Discussion

In the Harvest Management Plan for the Porcupine Caribou Herd in Canada (HMP; Porcupine Caribou Management Board 2010), there is a provision for bull only harvest to be implemented for different user groups if the herd drops below a certain population size. In addition, the PCMB continues to promote harvesting of bulls, regardless of population size. Population modeling has shown that if the proportion of bulls in the harvest rose from 30% to 80%, we could see a sex ratio in the herd of about 40 bulls per 100 cows. We don't really know what might happen to the herd sex ratio when we take more bulls during harvesting activities; as a result, we completed composition counts to get an updated bull ratio in 2009 and 2010 prior to the projected increase in harvested bulls resulting from the HMP.

The PCTC continues to recommend collecting this data occasionally, recognizing the very complex logistics and difficulties often experienced with weather during the rut period. A rut composition is most useful in calibrating population models for the herd. Accurate harvest data from all Parties, including the sex ratio of the harvest, is important to assess the effect of a bull dominated harvest on the herd sex ratio and to most accurately estimate the population size of the herd using models.

CARIBOU BODY CONDITION

Hunter assessments and condition indicators

Objective

This long term project uses specific samples from hunter killed caribou to track the fatness of Porcupine caribou.

Methods

Starting in 1987, Anne Allaye-Chan (a PhD student from University of Alaska Fairbanks) developed equations to estimate the body weight, body fat and body protein for adult cow Porcupine caribou (Allaye-Chan 1991). Government of Yukon (YTG) did collections from 1989 to 1991 to test these equations and in 1991, started regular monitoring with hunters from Old Crow (Porcupine River in September), Ft. McPherson, Dawson and Mayo (Yukon portion of the Dempster Highway in November and March).

In 2001, the PCTC formally modified the program so that hunters could submit samples from any caribou they harvest. This program is also called the Caribou Sampling Initiative (CSI) in the HMP and is also similar to the Circum-Arctic Rangifer Monitoring and Assessment Network Level 1 monitoring (Gunn and Nixon 2007). Hunters record a number of variables and rate the condition of their caribou.

Results

During the fall of 2020-21 caribou were generally not available to hunters. Small groups of caribou were initially available on the Dempster Highway and west of Aklavik, but by late August the majority of caribou had moved into Alaska. Caribou didn't stop moving however, and by mid to late September had reversed direction and returned to the Yukon. Caribou became available to hunters in Old Crow and from the Dempster Highway at the NWT border in late October. By November 2020 large numbers had also moved into the Richardson mountains and became available to hunters in Aklavik, NWT. The majority of samples for Body Condition analysis were collected by NWT hunters from November 2020 thru March 2021.

One hundred and sixty-six caribou were assessed by hunters for body condition, and 142 were measured for backfat. Overall, caribou condition in 2020 – 21 remained good, although slightly below the 5-year average. Backfat depths were below the overall average, but remained near the 5-year average.

Discussion

Average body condition data seems to be more variable after 2001 when hunters began rating their harvested caribou compared to when they were working with the biologists on the collection (Figure 8 and Figure 9). This could also be a seasonal effect; caribou collections in the early 1990's were done three times (Sept, Nov and March) whereas the current system allows hunters to submit samples all winter long. Bulls harvested in September tend to have the highest body condition and backfat values, while bulls and cows harvested in October and November tend to be significantly lower – as observed during this sampling period. Hunters also reported to biologists that cow caribou were thin in late November but by late winter had put on significant fat reserves, appearing to be in much better condition.

We should also keep in mind that hunters can be very selective when harvesting. This indicator gives an index of harvested caribou, not an index of the entire herd. Also, data are pooled over each winter but sample sizes remain small.




Figure 8. Average condition of harvested Porcupine caribou recorded by hunters. 1=poor 2=fair 3=good 4=very good. Error bars are standard errors. Labels indicate # of caribou sampled.




Figure 9. Average depth of backfat (cm) recorded in Body Condition Monitoring. Error bars are standard errors. Labels indicate # of caribou sampled.

HABITAT

Wildland fires

2021 season fire map data is not yet publicly available from Yukon or NWT. This section of the report contains information current to 2020 for Yukon and Alaska.

Objective

To monitor the amount of Porcupine Caribou range burned as an index of range condition.

Methods

Historical fire perimeter data was downloaded from the respective agencies websites. Some judgments were made to delete what we thought were duplicate fires and merge incompletely mapped fires along the borders between jurisdictions. Fire polygons were clipped to the extent of PCH range and total area burned was summed for each year. The Alaskan fire perimeter data starts in 1945, Yukon in 1945 and NWT in 1965, therefore only fire information between 1960-2020 was summarized in this report.

Results

As of the 2020 season, the total area burned by fires since 1960 is 46,778.3 km² or roughly 18% of the Porcupine Caribou herd's total annual range (Figure 10). **Areas burned within range of the Porcupine Caribou Herd in Yukon, NWT and Alaska from 1960 to 2020.** Yukon fires in 2020 burned a total of about 211.5 km², well below the 5-year average (1551.9 km²). In 2020 there were 6 fires in the Alaskan portion of the PCH range, 4 in the Yukon and none in the NWT (Figure 11). All of the fires recorded were considered small (<10,000 hectares). Although data is not available for 2021, only one major fire occurred in the core of the herd's range while several occurred on the periphery of their range.

Discussion

Fire perimeters are mapped by the fire management sections of the three jurisdictions. Although there are many similarities in methods, there are five cautionary notes when considering the data presented here. Firstly, the technology for remotely detecting wildland fires improved only in the 1960's therefore data prior to that should be viewed with caution. Secondly, past fires are continually being digitized from satellite or other remote sensing methods so the dataset will change as new data on old fires is added. Thirdly, maps show perimeters of fires only and do not reflect any unburned patches or varying fire severity within burned area. Fourthly, some fires are too small to map and are not included in the map files, and finally some fires burn areas that were previously burned.

There is much variability in how fires affect caribou; however, research completed on the Beverly Caribou Herd found that forests burned by wildfire produced enough lichen forage as early as 40 or 50 years after the fire, while research in central Yukon found some areas took over a century to recover. Caribou also tended to avoid burns larger than 10,000 hectares (100 km²). The rate of re-growth of caribou forage can be quite variable and caribou use of burns is generally unknown, therefore wildland fire information presented here should be considered as an index of changes to winter habitat.




Figure 10. Areas burned within range of the Porcupine Caribou Herd in Yukon, NWT and Alaska from 1960 to 2020.

Total number of fires and number of large fires
(greater than 10,000 hectares) in PCH range, by year




Figure 11. Total number of fires and number of large fires to 2020 within the range of the Porcupine Caribou Herd in Yukon, NWT and Alaska.

Total area of PCH range burned, by year




Figure 12. Total area burned by fire, by year to 2020 within the range of the Porcupine Caribou Herd in Yukon, NWT and Alaska.

Linear disturbance and human development footprint

Objective

To monitor the amount of linear disturbance and development footprint present on the herd's range.

Methods

Data is acquired from a number of sources in NWT, Yukon, and Alaska. Much of the historical data was acquired during a cumulative effects assessment completed for the PCMB in 2008-2012. Newer data was acquired for specific activities known to have been carried out within the range of the herd. Data quality varies for historical data but is thought to be more accurate for recent disturbances. Smaller developments (e.g., individual gravel quarries) may not be accounted for. In the case of historical disturbances (e.g., seismic lines cut in the 1960's), it is unknown whether features have adequately recovered or not to be removed from datasets, and in most cases a wide range of vegetation recovery can be expected even on the same feature. Recent work by the Yukon Land Use Planning Commission has evaluated historical disturbances and vegetation recovery in portions of the herd's range and this data may need to be evaluated in future years.

Data is mapped at a range scale and areas with significant new development are provided with a map of the specific area that has been disturbed. Where appropriate the total linear footprint has been summarized for different disturbance types and a total area of the footprint provided where possible.

Results

In 2020-21, no detectable changes occurred in linear disturbance and human footprint although some lease sales occurred in the 1002 area of the Arctic National Wildlife Refuge (ANWR). This area overlaps with PCH calving, post-calving and summer ranges. A relatively small (i.e., ~20 wells) is likely to be proposed for the Eagle Plains area in 2021, although that application remains forthcoming.

Most development in the range of the Porcupine Caribou herd occurred prior to the 1980's so we know relatively little about the disturbances except when they are still active (Figure 13). In 2013-2014, a major 3D seismic project occurred in the Eagle Plains area of Yukon. Based on data provided by Yukon Oil and Gas Branch and the company responsible for conducting the work, a total of 2,124 km of seismic line varying in width from 1.75-5 meters was cut, totaling approximately 5.35 km² of footprint. Access roads in the area totaled 228 km and varied in width between 3-5 meters (Figure 14).

Discussion

Linear disturbances and human footprint can affect caribou in multiple ways. Increased access can provide hunters with increased success and in some cases may facilitate predator movements resulting in higher predation levels or increased stress levels for caribou. Footprint can also impact habitat and habitat use by either directly impacting the habitat or by creating behavioural responses where caribou do not use high quality habitat as they may perceive it to be too risky. Large patches of intact habitat are known to be critical to caribou herds.

Snow condition

Objective

To gather an index of snow depth and hardness.

Methods

Water Resources (when under Environment Canada and now under Yukon Government) recorded late winter snow depth and snow water equivalent back to the 1970's. The Yukon Fish and Wildlife Branch also did late winter snow measurements along the Dempster Highway and Yukon north coast since the 1990's.

At specified permanent locations, ~10 repeated measures of depth and either snow density or snow water equivalent (SWE) are recorded. Where necessary, SWE is converted to density by dividing SWE by the depth of snow. Not all stations are measured in all years pending availability of staff. Data presented in this report represents results from 17 stations from the Yukon since 2013. Data from other jurisdictions were available in a compatible format for this report. For example the GNWT records SWE however does not record depth, so snow density measurements cannot be calculated and do not appear on Figure 16.

Results

The Porcupine Caribou Herd was spread out over portions of Alaska, Yukon and NWT during winter 2020 – 2021 (**Error! Reference source not found.**). In Alaska, caribou stretched between The Dalton Highway and Arctic Village, but few caribou wintered between Arctic Village and the Colleen River. Caribou were spread out east of the Colleen River thru the western edge of the Old Crow Flats and north into the upper Firth River. Caribou were spread out across the Yukon from the Alaska border into the upper Bluefish River, east up the Porcupine River, northeast up the Driftwood and Bell Rivers into the Richardson mountains, with a significant number of caribou wintering west of Alkavik. Caribou also extended southeast thru the Whitefish Wetlands into the upper Eagle River to the Dempster Highway.




Figure 15. Winter distribution of Porcupine Caribou from Dec. 1, 20120 to March 31, 2021.

Snow depth in the Eagle and Old Crow Regions were above the long term average, but below the 5 year average, while snow in the Ogilvie Region was above both the long and 5 year average. Snow density in all areas were slightly below the 5 year average, and below the long term average in all areas but Ogilvie (Figure 16).

Snow measures were not collected in the Richardson or North Slope regions. The long term trend for snow pack in the Eagle and Old Crow Regions appears to be increasing slightly, particularly over the last 10 years (Figure 17). Snow density may also be increasing in the Ogilvie and Old Crow Regions (Figure 18).

During late winter fieldwork in the Yukon and Alaska, we observed snow conditions that corresponded to the above data with us frequently observing snow conditions that appeared to be above average.

Discussion

When snow is deep or hardened by wind, caribou expend more energy digging through the snow which can potentially affect their body condition, and reproductive capability. Caribou are not always in the areas where we measure snow but this information can be used as an index of winter conditions affecting caribou.




Figure 16. Summary of snow depth and density by snow region from permanent stations (indicated by green stars) for the Yukon portion of the Porcupine Caribou Herd range. Red lines on the map delineate snow regions relevant to caribou (Russell et al 1993).

LITERATURE CITED

- Allaye-Chan, A.C. 1991. Physiological and ecological determinants of nutrient partitioning in caribou and reindeer. Ph.D. Thesis. University of Alaska Fairbanks, AK. 125 pp.
- Arthur, S. 2001. Alaska Department of Fish and Game memo dated 13 November 2001.
- Arthur, S. M., K. R. Whitten, F. J. Mauer and D. Cooley. 2003. Modeling the decline of the Porcupine Caribou Herd, 1989-1998: the importance of survival vs. recruitment. *Rangifer*. Special Issue 14:123-130.
- Boertje, R. D., C. L. Garder, K.A. Kellie, and B.D. Taras. 2012. Fortymile caribou herd: Increasing numbers, declining nutrition and expanding range. Alaska Department of Fish and Game, Wildlife Technical Bulletin 14, ADF7G/DWC/WTB-2012-14. Juneau, Alaska.
- Caikoski, J.R. 2017. Photocensus of the Porcupine Caribou Herd. Alaska Department of Fish and Game, Division of Wildlife Conservation. Memorandum, December 20, 2017. Fairbanks, Alaska.
- Caikoski, J. R. 2018. Porcupine Caribou management report and plan, Game Management Unit 25A, 25B, 25D, and 26C: Report period 1 July 2012–30 June 2017, and plan period 1 July 2017–30 June 2022. Alaska Department of Fish and Game, Species Management Report and Plan ADF&G/DWC/SMR&P-*in press*, Juneau.
- Caikoski, J.R. 2020a. Porcupine Caribou Herd Calving and Post Calving Surveys, May-June 2020. Alaska, Department of Fish and Game, Division of Wildlife Conservation. Memorandum, July 2020. Fairbanks, Alaska.
- Caikoski, J. R. 2020b. Porcupine caribou herd management report and plan, Game Management Unit 25A, 25B, 25D, and 26C: Report period 1 July 2012–30 June 2017, and plan period 1 July 2017–30 June 2022. Alaska Department of Fish and Game, Species Management Report and Plan ADF&G/DWC/SMR&P-2020-22, Juneau.
- Caikoski, J.R. 2021. Porcupine Caribou Herd Calving and Post Calving Surveys, May-June 2021. Alaska, Department of Fish and Game, Division of Wildlife Conservation. Memorandum, August 2021. Fairbanks, Alaska.
- Davis, J. L., P. Valkenburg, and S. Harbo. 1979. Refinement of the aerial photo-direct count-extrapolation caribou census technique. Alaska Department of Fish and Game. Federal Aid in Wildlife Restoration, Research Progress Report, Grant W-17-11, Study 3.25, Juneau.
- Fancy, S. G, K. R. Whitten, and D. E. Russell. 1994. Demography of the Porcupine caribou herd, 1983–1992. *Canadian Journal of Zoology* 72:840–846.
- Griffith, B., D. C. Douglas, N. E. Walsh, D. D. Young, T. R. McCabe, D. E. Russell, R. G. White, R. D. Cameron, and K. R. Whitten. 2002. The Porcupine caribou herd. Pages 8–37 [In] D. C. Douglas, P. E. Reynolds, and E. B. Rhode, editors. Arctic Refuge coastal plain terrestrial wildlife research summaries. U.S. Geological Survey, Biological Resources Division, Biological Science Report USGS/BRD/BSR-2002-0001. Reston, Virginia.

- Gunn, A. and W. Nixon. 2007. Rangifer Health & Body Condition Monitoring Manual. Circum Arctic Rangifer Monitoring and Assessment (CARMA) Network.
- International Porcupine Caribou Board. 1993. Plan for the International Conservation of the Porcupine Caribou Herd. Appendix ii of The International Porcupine Caribou Board – Fourth Annual Report 1995. 20 pp.
- Kaplan, E. L. and P. Meier. 1958. Nonparametric estimation from incomplete observations. *Journal of American Statistics Association* 53:457–481.
- Lenart, E.A. 2007. Game Management Units 25A, 25B, 25D and 26C. Pages 232 – 248. In P. Harper, editor. Caribou management report of survey-inventory activities, 1 July 2004 to 30 June 2006. Alaska Department of Fish and Game. Project 3.0. Juneau, Alaska, USA.
- Pollock, K. H., S. R. Winterstein, C. M. Bunck, and P.D. Curtis. 1989. Survival analysis in telemetry studies; the staggered entry design. *Journal of Wildlife Management* 53:7–15
- Porcupine Caribou Management Board. 2010. Harvest Management Plan for the Porcupine Caribou Herd in Canada. March 2010. 45 pp.
- Porcupine Caribou Management Board. 2015. Porcupine Caribou Herd Strategic Framework 2015-16 to 2019-20. 12 pp.
- Rivest, L-P., S. Couturier, and H. Crepeau. 1998. Statistical methods for estimating caribou abundance using postcalving aggregations detected by radio telemetry. *Biometrics* 54:865–876.
- Russell, D.E., A.M. Martell and W.A.C. Nixon. 1993. Range ecology of the Porcupine Caribou Herd in Canada. *Rangifer Special Issue No 8*. 168 pp.
- Russell, D.E., M.Y. Svoboda, Arokium J. ,Cooley D. 2013. Arctic Borderlands Ecological Knowledge Cooperative: can local knowledge inform caribou management? *Rangifer* (33) Special Issue No 21 71-78.
- Thomas, D. C. and H. P.L. Kiliaan. 1998. Fire-caribou relationships: (IV) Recover of habitat after fire on the winter range of the Beverly Herd. Technical Report Series No. 312. Canadian Wildlife Service, Environment Canada, Prairie and Northern Region.
- Valkenburg, P., D. A. Anderson, J. L. Davis, and D. J. Reed. 1985. Evaluation of an aerial census technique for caribou based on radiotelemetry. Pages 287–299 [In] T. C. Meredith and A. M. Martell, editors. Proceedings second North American caribou workshop. McGill Subarctic Research Paper 40.
- Wertz, T.L., S.M. Arthur, D. Cooley, B. Griffith, and M. Kienzler. 2007a. Seasonal Survival of The Porcupine Caribou Herd In Alaska And Northern Yukon Territory, 2003-2006. Project report to the Porcupine Caribou Management Board.
- Wertz, T. L., S. M. Arthur, B. Griffith, D. Cooley, M. Kienzler. 2007b. The puzzle of the Porcupine caribou herd: Do we have all the pieces? Yukon North Slope Conference. Whitehorse, Yukon.
- Whitten, K. R. 1995. Antler loss and udder distention in relation to parturition in caribou. *Journal of Wildlife Management* 59:273–277.

Appendix A. Summary of biological parameters

Year	Parturition Rate ≥ 4 yrs (sample size)	Parturition Rate ≥ 3 yrs (sample size)	Parturition Rate ≥ 3 yrs (sample size)	June Calf Survival ^c	Post-calving Survival ^d	Late June Calf: Cow ^e	March Calf: Cow ^f	Population Estimate	Peak of calving	Calving note
1985	0.77					0.46				
1986	0.74									
1987	0.78 (51)			0.71		0.55		165,000		
1988	0.84 (91)			0.65		0.55				
1989	0.78 (74)			0.74		0.58	0.43	178,000		
1990	0.82 (74)			0.90		0.74				
1991	0.74 (77)			0.82		0.61	0.22			
1992	0.86 (78)			0.57		0.49	0.33	160,000		
1993	0.81 (63)			0.56	0.83	0.45	0.32			
1994	0.91 (98)			0.77	0.93	0.70	0.40	152,000		
1995	0.69 (95)			0.85	0.92	0.59	0.41			
1996	0.89 (74)			0.81	0.91	0.72	0.46			
1997	0.75 (48)			0.77	0.90	0.58	0.38			
1998	0.83 (58)			0.82	0.94	0.68	0.27	129,000		
1999	0.84 (39)			0.83	0.86	0.70	0.56		3-Jun	1-5 June
2000	0.73 (44)			0.61	0.82	0.44	0.28		7-Jun	
2001	0.84 (70)			0.61	0.79	0.51	0.31	123,000	8-Jun	5-10 June
2002	0.87 (68)			0.65	0.85	0.56	0.38		5-Jun	
2003	0.87 (70)			0.79	0.85	0.69	0.33		1-Jun	
2004	0.82 (74)			g	g	g	0.24		3-Jun	3-4 June
2005	0.64 (55)	0.63 (65)	0.60 (10)	0.77	0.88	0.49	h		2-Jun	1 - 4 June
2006	0.79 (66)	0.79 (67)	1.00 (1)	0.73	0.86	0.58	0.39		2-Jun	
2007	0.88 (67)	0.89 (71)	1.00 (4)	0.83	0.90	0.73	h		30-May	
2008	0.79 (63)	0.79 (69)	0.83 (6)	0.73	0.92	0.59	h		30-May	29 or 30 May
2009	0.77 (65)	0.79 (72)	1.00 (7)	0.57	0.75	0.44	0.19		2-Jun	

Year	Parturition Rate ≥ 4 yrs (sample size)	Parturition Rate ≥ 3 yrs (sample size)	Parturition Rate ≥ 3 yrs (sample size)	June Calf Survival ^c	Post-calving Survival ^d	Late June Calf:Cow ^e	March Calf:Cow ^f	Population Estimate	Peak of calving	Calving note
2010	0.85 (41)	0.75 (48)	0.14 (7)	0.76	0.87	0.65	h	169,000	1-Jun	prior to 2 Jun
2011	0.86 (59)	0.86 (59)		0.48	0.59	0.41	h		30-May	prior to 1 Jun
2012	g	g	g	g	g	g	g		30-May	prior to 1 Jun
2013	0.86 (42)	0.85 (45)	0.67 (3)	i	l	i		197,000	4-Jun	3-4 June
2014	g	g	g	g	g	0.49			no data	
2015	g	g	g	g	l	i			no data	
2016	0.75 (28)	0.76 (37)	0.78 (9)	0.61	1.00	0.46	0.36		3-Jun	
2017	0.90 (42)	0.92 (54)	1.00 (12)	0.81	0.90	0.72		218,000	3-Jun	
2018	0.88 (41)	0.78 (50)	0.33 (9)	0.73 ^j	0.88 ^j	0.64 ^j			6-Jun	
2019	0.81 (42)	0.72 (57)	0.47 (15)		0.94 ^j	0.56 ^j			4-Jun	
2020	0.80 (35)	0.79 (42)	0.71 (7)		0.93 ^j	0.57 ^j			1-Jun	
2021	0.85 (39)	0.83 (46)	0.71 (7)		0.90 ^j	0.63 ^j			30-May	
Mean	0.82	0.79	0.60	0.72	0.87	0.57	0.35		31-May	
5 yr mean	0.85	0.81	0.64	0.72	0.91	0.58	0.36		2-Jun	

^a Data are from Fancy et al. (1994, Can. J. Zool. 72:840–846), Alaska Department of Fish and Game, and Yukon Department of Environment.

^b Number of radiocollared adult cows for which parturition status was determined in early June, excluding those known to be <4 years old. Includes caribou of unknown age, but most likely > or equal to 4 years olds. Prior to 2003, all caribou were of unknown age.

^c Estimated as (July calf:cow ratio)/(parturition rate).

^d Includes only calves observed during early June that were subsequently observed in late June (i.e., does not include most perinatal mortality).

^e Excludes radiocollared cows known to be < 4 years old.

^f As of March of the year following birth of each cohort; includes all cows >1 year old.

^g No data due to adverse weather conditions.

^h No data due to mixing of herds on winter range.

ⁱ No data due to dense caribou groups making identification of cow:calf pairs not possible.

^j Starting in 2018, all females ≥3 years old are included in the summary. Past status reports only reported adult cows ≥4 years of age unless otherwise stated in the footnotes. Parturition status remains for radiocollared adults ≥4 yrs old.

Appendix B. Previous research findings

Adult female survival

Objective

To obtain an annual estimate of survival for adult female Porcupine caribou.

Methods

There have been a number of issues with past methods to determine female survival estimates. However, with increases in the number of GPS collars deployed on the herd, the PCTC can calculate survival estimates using the known fate (i.e. alive, dead) of GPS collared females. Analyses will include how survival varies by age, sex and season.

Results

Researchers started a project in 2003 to get an updated estimate of adult female survival in response to the continued population decline (Wertz et al 2007a). Survival estimates ranged between 0.065 to 0.097 but showed a general trend of increased survival from 2003 to 2011. **(Figure 17. Annual survival estimates for adult female Porcupine Caribou, May 2003 – June 2012. Source: USFWS unpublished data.)**

Discussion

As with many populations, the survival of breeding females is very important to the potential growth of the herd. A sustained change of 2 or 3 percent in survival can make the difference between a herd increasing and decreasing. Adult female survival has been estimated twice before; once when the herd was increasing and again when the herd began to decline (Fancy et al 1994, Walsh et al 1995). Information gathered from these earlier studies indicated that most cows died in winter, the harshest season of the year.




Figure 17. Annual survival estimates for adult female Porcupine Caribou, May 2003 – June 2012. Source: USFWS unpublished data.

Short yearling survival to 3 years of age

Objective

To document the survival of 9 month old calves to 3 years of age (2003-2010 only).

Methods

Starting in 2003, we captured about 10 female caribou in March that were born the previous spring (9 months old) and put conventional radio collars on them. The data from all years of captures were pooled to estimate how many calves survived to breeding age. Because we know exactly how old these caribou were, we recapture them after 3 years or sooner and replaced their collars to maintain a collared sample.

Results

The average survival rate of female Porcupine caribou appears to decline as caribou age from 9 months to 3 years but because of the error bars overlap on the estimates we cannot say for certain that there is any change in survival rate as caribou survive to breeding age (**Figure 18. Survival of Porcupine Caribou females from 9 months to 3 years of age from 2003-2010.**). The average survival rates of female caribou 9 mo. – 3 yrs of age are similar (87%) to adult female survival rates taken from the same time period (84%). The last year of this seven year study was completed in 2010.

Discussion

In 2003, we started a 7-year study to estimate how well calves survive to 3 years of age when they should enter the breeding portion of the population. This has been estimated only once before in 1983-88 (Fancy et al 1994). We have been assuming that once calves reach one year of age, they survive at the same rate as adults. We are testing this assumption because, as with the survival of adult females, the survival of young females is important to population dynamics. Computer population modeling shows that it would take a decrease of only 6% in adult female survival or a decrease of 50% of calves to cause a decline like we have documented for the Porcupine Caribou Herd between 1989 and 2001 (Arthur et al 2003). Other work has shown that survival of calves in their first year of life is very low. Survival of these young, non-breeding animals is similar to adult females.

Small sample sizes are an issue for this analysis. The estimates are based on data pooled over multiple years of collaring efforts, however the sample size at step one of the analysis is 59 animals. In order to be able to detect small changes in short yearling survival with confidence, we would have to maintain collars on many more young caribou. There are constraints to doing that in terms of funding, availability of free radio frequencies, logistics of flying, and community concerns. Despite these constraints, we decided to continue small numbers collaring short yearling females each year to continue recording survival estimates (low statistical power given the small sample size) but also to ensure the collared sample of caribou is not biased toward older animals.




Figure 18. Survival of Porcupine Caribou females from 9 months to 3 years of age from 2003-2010.

Adult bull survival

Objective

To document the survival of adult bull caribou (2003-2010 only).

Methods

Each year before a census attempt, we deploy a number of collars on adult bull caribou so we can locate the bull groups during the census field work. Because we've been preparing for a census each year for 8 years running, we have an unprecedented number of bulls collared. We are able to do an analysis similar to the short yearling analysis. All collared bulls were pooled and we calculated their survival rate in years following capture.

Results

Between 2003 and 2006, more bulls died during the fall than any other season. Bull mortality rate increases dramatically about 5 years after collaring (**Figure 19. Survival of male Porcupine Caribou from 2003 to 2010.**). Assuming bulls were at least 3 years old at the time of capture, bulls start dying at an increased rate at 8 or more years of age. The study on adult bull survival extended from 2003 – 2010. No further collaring of bulls is planned.

Discussion

As expected, we see that bulls seems to survive at a lower rate than adult cows. Bulls are probably more stressed during the rut which contributes to a lower survival rate.




Figure 19. Survival of male Porcupine Caribou from 2003 to 2010.